

DEVELOPMENT OF A MULTIPLEX PCR METHOD FOR DETECTION OF THE GENES ENCODING 16S rRNA, COAGULASE, METHICILLIN RESISTANCE AND ENTEROTOXINS IN *STAPHYLOCOCCUS AUREUS*

YILING FAN¹, FENG PAN¹, GEORGE C. PAOLI², YONGHONG XIAO³,
HAIHUI SHENG⁴ and XIANMING SHI^{1,5}

¹*Department of Food Science and Bor Luh Food Safety Center
School of Agriculture and Biology
Shanghai Jiao Tong University
800 Dong Chuan Road, Shanghai 200240, China*

²*Microbial Biophysics and Residue Chemistry
Eastern Regional Research Center
Agricultural Research Service
United States Department of Agriculture
600 East Mermaid Lane, Wyndmoor, PA*

³*Institute of Clinical Pharmacology
Peking University
Beijing, China*

⁴*Shanghai Biochip Co., Ltd.
Shanghai, China*

Accepted for Publication April 10, 2008

ABSTRACT

A multiplex polymerase chain reaction (PCR) method was developed for simultaneous detection of the genes encoding methicillin resistance (mecA), SEs A, B and C (sea, seb and sec), coagulase (coa), and 16S rRNA. The primers for amplification of the 16S rRNA gene were specific for Staphylococcus spp., and the primers for coa were specific for Staphylococcus aureus. Based on the results, the multiplex PCR was accomplished at an optimal Mg²⁺ concentration of 1.0 mM and at an annealing temperature of 56°C. This multiplex PCR method was performed with 71 strains of S. aureus and 51 strains of six other bacterial species. Among the S. aureus strains tested, 40.0% (28/71) were found to contain the mecA gene. One of the 28 mecA⁺ strains was not resistant to methicillin. The sea, seb and sec genes were present in 47.9%

⁵Corresponding author. TEL: 86-21-3420-6616; FAX: 86-21-3420-6616; EMAIL: xmshi@sjtu.edu.cn

(34/71), 5.6% (4/71) and 8.5% (6/71) of *S. aureus* strains, respectively. The sensitivity of this multiplex PCR method was approximately 104.5 pg of genomic DNA per reaction, which was equivalent to an estimated 2.4×10^3 cfu of *S. aureus* or 3.64×10^4 copies of genome equivalent.

PRACTICAL APPLICATIONS

The developed multiplex polymerase chain reaction (PCR) method will be a useful tool for the detection and identification of *Staphylococcus aureus* from foods, clinical samples and environmental surveys. In particular, the routine use of this multiplex PCR method for the detection of foodborne *S. aureus* could be used to monitor the presence of enterotoxins and the emergence of methicillin resistance in a population that, to date, has had a relatively low incidence of methicillin resistance.

INTRODUCTION

Staphylococcus aureus, a gram-positive bacterium, is one of the most common clinical and foodborne pathogens worldwide (Lowy 1998; Weigel *et al.* 2003; Foster 2005; Cagatay *et al.* 2007). *S. aureus* produces many important virulence factors including SEs, which are the main causes of diarrhea, vomiting and other symptoms associated with *S. aureus* infection. It was reported that more than 70% of *S. aureus* strains produced one or more enterotoxins (Jorgensen *et al.* 2005). Because of their thermal stability, SEs remain a great hazard even in heat-processed foods (Akineden *et al.* 2001). Several SEs from SEA to SEE, SEG to SEO, SEU and SEV have been characterized according to serological analysis and classification, and most of their genes have been sequenced (Akineden *et al.* 2001; Omoe *et al.* 2002; Thomas *et al.* 2006; Vimercati *et al.* 2006). Five of these SEs (SEA to SEE) are recognized as major causes of foodborne illness (Vimercati *et al.* 2006), with SEA, SEB and SEC being the most frequently implicated in foodborne outbreaks (Balaban and Rasooly 2000).

There has been an enormous increase in the isolation of MRSA strains that has been attributed to the widespread use of methicillin in clinical settings and in food animal production facilities for decades (Panlilio *et al.* 1992; Oliveira *et al.* 2002). In addition, the gene encoding methicillin resistance (*mecA*) has been widely used as a molecular marker of methicillin resistance in detection and typing of *Staphylococcus* spp. (Ubukata *et al.* 1990; Carroll *et al.* 1996; Jonas *et al.* 2002; Francois *et al.* 2003). SEs increase the competi-

tiveness of *S. aureus*, and antibiotic resistance gives *S. aureus* a greater ability to distribute widely (Massey *et al.* 2006).

Recently, many polymerase chain reaction (PCR) methods have been developed to detect enterotoxin genes or methicillin resistance-related genes of *S. aureus* (McLauchlin *et al.* 2000; Mehrotra *et al.* 2000; Jonas *et al.* 2002; Francois *et al.* 2003; Padmapriya *et al.* 2003). Some of those methods were developed to detect several *S. aureus* toxins and methicillin resistance, but the SEs and methicillin resistance genes were not amplified in the same reaction (Mehrotra *et al.* 2000). To our knowledge, the identification of both SEs and methicillin resistance genes in a single multiplex PCR has not been reported.

The 16S rRNA gene of staphylococci contains DNA sequences that are highly conserved at the genus level but are variable among other bacterial genera (Baron *et al.* 2004). Several primers designed against the 16S rRNA gene have proven useful for identification of staphylococci (Mason *et al.* 2001; Yang *et al.* 2002). The 16S rRNA gene primer pair used in this multiplex PCR method was previously shown to be conserved in 19 *Staphylococcus* species and subspecies (Zhang *et al.* 2004). In addition, the coagulase gene (*coa*) was reported as a specific target for the detection of *S. aureus* at the species level (Chiou *et al.* 2000; Karahan and Cetinkaya 2006; Sabat *et al.* 2006). Therefore, the combined detection of both the 16S rRNA and coagulase genes may be a reliable set of markers for the detection of *S. aureus*.

The aim of this study was to identify *S. aureus* strains via amplification of the genes encoding the 16S rRNA and coagulase, and to determine the presence of the methicillin resistance gene (*mecA*) and the genes encoding the most common foodborne outbreak-associated SEs (*sea*, *seb* and *sec*) in a single multiplex PCR.

MATERIALS AND METHODS

Strains and Culture Medium

A total of 135 bacterial strains (Table 1) were cultured in Luria–Bertani broth at 37°C for 6 h with shaking at 250 rpm before extraction of genomic DNA. All staphylococci isolates were separated from clinic samples and were identified by the Institute of Clinical Pharmacology at Peking University using API Staph Ident (bioMérieux, Shanghai, China).

Identification of Methicillin Resistance

The methicillin resistance of *S. aureus* strains was tested by the standardized agar dilution method (Wikler *et al.* 2006). The experiments were

TABLE 1.
BACTERIAL STRAINS USED IN THIS STUDY

Bacterial species/strains	Number	Source
<i>Staphylococcus aureus</i> isolates	67	*
<i>S. aureus</i> ATCC6538	1	†
<i>S. aureus</i> ATCC25923	1	†
<i>S. aureus</i> ATCC29213	1	†
<i>S. aureus</i> CMCC26001	1	†
<i>S. aureus</i> ATCC 27940	1	‡
<i>S. aureus</i> ATCC 13565	1	‡
<i>Staphylococcus haemolyticus</i> strains	8	*
<i>Staphylococcus epidermidis</i> strains	7	*‡
<i>S. saprophyticus</i> ATCC 15305	1	‡
<i>S. saprophyticus</i> ATCC 43809	1	‡
<i>S. lugdunensis</i> ATCC 43809	1	‡
<i>S. schleiferi</i> ATCC 43808	1	‡
<i>S. xylosus</i> ATCC 29971	1	‡
<i>S. sciuri</i> ATCC 29062	1	‡
<i>S. capitis</i> ATCC 49325	1	‡
<i>S. chromogenes</i> 3688	1	‡
<i>S. simulans</i> 13044	1	‡
<i>S. intermedius</i> ATCC 29663	1	‡
<i>Salmonella</i> spp. strains	5	†
<i>Escherichia coli</i> strains	22	†
<i>Listeria monocytogenes</i> strains	4	†
<i>Vibrio parahaemolyticus</i> strains	6	†
Total	135	

*Clinical isolates were obtained from Peking University Health Science Center, Beijing, China.

†Bacterial strains were acquired from Bor Luh Food Safety Center, Shanghai Jiao Tong University, Shanghai, China.

‡Dr. George C. Paoli, Microbial Biophysics and Residue Chemistry, Eastern Regional Research Center, Agricultural Research Service, United States Department of Agriculture, Wyndmoor, PA, USA.

carried out at the Institute of Clinical Pharmacology at Peking University. Briefly, *S. aureus* colonies were taken from an overnight sheep blood agar plate, and a suspension was prepared in Mueller–Hinton broth equivalent to that of the 0.5 McFarland standards. Approximately 10^4 cfu of each isolate was inoculated on Mueller–Hinton agar (supplemented with 4% NaCl) containing serial twofold dilutions of methicillin (16, 8, 4, 2 or 1 $\mu\text{g/mL}$). The microdilution trays were incubated at 35C for 24 h. Strains were identified as methicillin resistant in this study if the minimal inhibitory concentration was greater than 4 $\mu\text{g/mL}$ on agar plates after 24 h of incubation at 35C.

Genomic DNA Extraction

A modified cetyltrimethylammonium bromide method was used for extraction of bacterial genomic DNA (Kalia *et al.* 1999; Tang *et al.* 2006). One milliliter of an overnight culture was used to extract genomic DNA. The genomic DNA pellet was resuspended finally in 20 μ L sterile water and was stored at -20°C . The concentrations of the genomic DNA were determined by measuring the absorbance at 260 nm using a DU 800 UV/Visible Spectrophotometers System (Beckman, Shanghai, China).

PCR Primers

All primers in this study (Table 2) were analyzed by Premier v5.0 (PREMIER Biosoft International, Palo Alto, CA) and were synthesized by Shanghai Bio-tech Corporation (Shanghai, China).

DNA Amplification

One microliter of the bacterial DNA sample was added to 24 μ L PCR mixture containing 1 U of *rTaq* DNA polymerase (Tiangen Biotech Co. Ltd., Beijing, China); 1 \times Taq buffer (200 mM Tris-HCl [pH 8.4], 200 mM KCl and 100 mM $[\text{NH}_4]_2\text{SO}_4$); 1.0 mM MgCl_2 ; 0.25 mM of each deoxyribonucleotide triphosphate (Tiangen Biotech Co. Ltd.); 50 nM of each primer for the amplification of the 16S rRNA, *coa* and *mecA* genes; 100 nM of each primer for the amplification of *sea* gene; and 75 nM of each primer for the amplification of *seb* and *sec* genes. All PCR amplifications were carried out in Peltier Thermal Cycler PTC-200 (Bio-Rad Laboratories, Hercules, CA). The amplification conditions were as follows: initial denaturation at 94°C for 5 min; followed by 30 cycles of 94°C for 30 s, 56°C for 30 s and 72°C for 30 s; with a final extension at 72°C for 5 min before 16°C for 10 min.

Six microliters of the amplified PCR products was used for the separation by electrophoresis in a 2.0% agarose gel at 140 V in 1 \times Tris-acetate-ethylenediaminetetraacetic acid buffer for 40 min and visualization with UV illumination (Shanghai Tanon Science & Technology Co. Ltd., Shanghai, China) after staining with ethidium bromide for 15 min. Standard DNA molecular weight markers (100 bp or 1 kb DNA ladder, TianWei Science & Technology Co. Ltd., Beijing, China) were included in each run.

Optimization of Amplification Conditions

Each pair of primers was used in separate amplification reactions employing the protocol described previously. Three *S. aureus* strains were used for the optimization: *S. aureus* 05B038 (*mecA*⁺ and *sea*⁺), *S. aureus* 05L198 (*seb*⁺) and

TABLE 2.
PRIMER SEQUENCES USED IN THIS STUDY

Target gene	Primer	Oligonucleotide sequences (5'-3')	GenBank accession number	Amplicon size (bp)	Reference
16S rRNA	16S-F	AACTCTGTTATTAGGGAAGAACA	NC_002758.2	756	Zhang <i>et al.</i> 2004
	16S-R	CCACCTTCCTCCGGTTTGTCAAC			
<i>coa</i>	COA-F	CCTCAAGCAACTGAACAACA	BA000018.3	151	This study
	COA-R	TGAATCTTGGTCTCGCTTCAT			
<i>mecA</i>	MECA-F	GTAGAAATGACTGAACGTCCGATAA	X52593	310	Perez-Roth <i>et al.</i> 2001
	MECA-R	CCAATCCACATTGTTTCGGTCTAA			
<i>sea</i>	SEA-F	ATTAAACCGAAGGTCTGTAGA	M18970.1	552	Tang <i>et al.</i> 2006
	SEA-R	TTGGGTAAAAAGTCTGAATT			
<i>seb</i>	SEB-F	TGTATGTATGGAGGTGAAC	NC_002951.2	270	Sharma <i>et al.</i> 2000
	SEB-R	ATA GTG ACG AGTAG GTA			
<i>sec</i>	SEC-F	ACCAGACCCCTATGCCAGATG	AB084256	371	Cremonesi <i>et al.</i> 2005
	SEC-R	TCCCATTATCAAAAGTGTTTCC			

S. aureus SA6 (*sec*⁺). The gradient of annealing temperature ranged from 50.0 to 62.0°C, and the gradient of Mg²⁺ concentration ranged from 0.5 to 2.5 mM.

Specificity and Reliability Test

Seventy-one *S. aureus* strains and 51 strains of six other bacterial species were tested at the optimal multiplex PCR condition using the genomic DNA from *S. aureus* ATCC29213 strain (*sea*⁺) as a positive control, and double distilled water as a negative control. This multiplex PCR assay was further evaluated using 30.0 ng of the genomic DNA from the *S. aureus* strain 05B038 (*mecA*⁺ and *sea*⁺) in the presence of an equal amount of DNA from strains of *Salmonella*, *Escherichia coli*, *Listeria monocytogenes* or *Vibrio parahaemolyticus*.

Sensitivity Test

All target genes were examined in a series of multiplex PCRs to determine the sensitivity of the assay using the following strains: *S. aureus* 05B038 (*mecA*⁺ and *sea*⁺), *S. aureus* 05L198 (*seb*⁺) and *S. aureus* SA6 (*sec*⁺). These tests were carried out using the optimal Mg²⁺ concentration (1.0 mM) with 1 U of *rTaq* DNA polymerase at 56°C. The *S. aureus* DNA samples were prepared in a 10-fold dilution, and 1 µL of each dilution was used in this multiplex PCR assay. The quantities of the genomic DNA from the *S. aureus* strain 05B038 ranged from 1.045 to 1.045 × 10⁴ pg per reaction; the quantities of the genomic DNA from the *S. aureus* strain 05L198 ranged from 11.05 to 1.105 × 10⁵ pg per reaction; and the quantities of the genomic DNA from the *S. aureus* SA6 ranged from 2.757 to 2.757 × 10⁴ pg per reaction.

RESULTS

Optimization of Amplification Conditions

The primers for the amplification of the 16S rRNA, *coa*, *mecA* and *sea* genes worked well at all temperatures tested (details not shown). The annealing temperature of the multiplex PCR in this study was chosen at 56°C, and the PCR amplification products from each of the target genes at 56°C are shown in Fig. 1 (lanes 1–6).

When testing various Mg²⁺ concentrations, the product of the 16S rRNA gene gave a clearer and brighter band at 1.0 mM of Mg²⁺ (Fig. 2, lanes 2, 8 and 13) than at other Mg²⁺ concentrations (0.5, 1.5, 2.0 or 2.5 mM), and all the expected target genes were amplified at this concentration. Higher Mg²⁺ concentrations (more than 2.0 mM) might have inhibited the *rTaq* polymerase




FIG. 1. SINGLEPLEX AND MULTIPLEX POLYMERASE CHAIN REACTION (PCR) ASSAYS FOR THE DETECTION OF METHICILLIN RESISTANCE AND ENTEROTOXIN GENES IN *STAPHYLOCOCCUS AUREUS* STRAINS

Lane 1, 16S rRNA amplicon from *Staphylococcus aureus* 05B038; lane 2, *coa* amplicon from *S. aureus* 05B038; lane 3, *mecA* amplicon from *S. aureus* 05B038; lane 4, *sea* amplicon from *S. aureus* 05B038; lane 5, *seb* amplicon from *S. aureus* 05L198; lane 6, *sec* amplicon from *S. aureus* SA6; lanes 7–9, multiplex PCR products from *S. aureus* strains 05B038, 05L198 and SA6, respectively; lane 10, multiplex PCR using *Escherichia coli* genomic DNA; lane 11, 100 bp DNA marker.




FIG. 2. EFFECT OF Mg^{2+} CONCENTRATION ON THE MULTIPLEX POLYMERASE CHAIN REACTION (PCR) ASSAY

The concentrations of Mg^{2+} were 0.5, 1.0, 1.5, 2.0 and 2.5 mM from left to right for each strain. Lanes 1–5, *Staphylococcus aureus* 05B038; lanes 7–11, *S. aureus* 05L198; lanes 13–17, *S. aureus* SA6; lanes 6 and 12, 100 bp DNA marker.

activity, resulting in a reduced amplification in multiplex PCR (Fig. 2). Therefore, 1.0 mM Mg^{2+} was used in subsequent PCR experiments. The final condition of the multiplex PCR system, as determined from the optimization experiments, was 1.0 mM Mg^{2+} at an annealing temperature of 56°C for 30 cycles. As shown in Fig. 1, the tested primers of multiplex PCR worked efficiently at the chosen multiplex PCR conditions.

Specificity and Reliability Tests

A total number of 122 bacterial strains (Table 1) were tested using the multiplex PCR method. The *coa* and 16S rRNA genes were amplified from all *S. aureus* strains tested by this multiplex PCR (Table 3). To substantiate the multiplex PCR method, all *S. aureus* strains tested by the multiplex PCR were also screened for the presence of individual enterotoxin genes by singleplex PCRs. The results of the singleplex PCRs corresponded precisely with those from multiplex PCR (data not shown), demonstrating the reliability of the multiplex PCR method.

Amplicons were generated using the primers of the 16S rRNA gene, but not for the *coa* gene, when DNA from other species of *Staphylococcus* were tested using the multiplex PCR assay (Fig. 3). None of the 37 strains of other bacterial genera yielded any PCR products from any of the primer sets included in the multiplex PCR (data not shown). It is therefore concluded that the primers for amplification of the 16S rRNA gene were specific for *Staphylococcus* spp., as previously reported (Zhang *et al.* 2004). The specificity of the *coa* primers for the detection of *S. aureus* was also demonstrated by the results of the multiplex PCR tests described previously. The specificity of the *coa* primer set was further tested against additional strains of *Staphylococcus* spp. (Table 1 strains from source †) in a singleplex assay. The two additional strains of *S. aureus* yielded a *coa* PCR product, but none of the 11 other *Staphylococcus* spp., including the coagulase-positive species *Staphylococcus intermedius* and *Staphylococcus schleiferi*, yielded a *coa* amplicon (data not shown). The results of the multiplex assays (Table 3 and Fig. 3) and the singleplex PCR using the *coa* primers (data not shown) demonstrated the specificity of this primer set for *S. aureus*.

In addition, it was found from the tests by the multiplex PCR method that the *sea*, *seb* and *sec* genes were present in 47.9% (34/71), 5.6% (4/71) and 8.5% (6/71) of the tested *S. aureus* strains, respectively. In this study, 38.0% (27/71) of the tested *S. aureus* strains were both enterotoxin gene(s) (*sea*, *seb* or *sec*) and *mecA* negative; 8.5% (6/71) of the *S. aureus* strains were *mecA*⁺ and enterotoxin gene negative; and 29.6% (21/71) of the strains were *mecA*⁺ and had one or more enterotoxin gene(s). None of the MRSA strains yielded a *seb* amplicon. There were 28 *S. aureus* strains that yielded an *mecA* amplicon, but only 27 *S. aureus* strains were phenotypically resistant to methicillin. The *mecA* gene may not be properly expressed in the *mecA*⁺ methicillin-sensitive strain. The two strains of *Staphylococcus haemolyticus* and one strain of *Staphylococcus epidermidis* tested yielded 16S rRNA gene and *mecA* gene amplicons, but no enterotoxin genes were detected in these two *Staphylococcus* species (Fig. 3).

TABLE 3.
METHICILLIN RESISTANCE AND MULTIPLEX POLYMERASE CHAIN REACTION
RESULTS FOR *STAPHYLOCOCCUS AUREUS* STRAINS

<i>S. aureus</i> strain	Gene amplified						Methicillin*	Total
	<i>sea</i>	<i>seb</i>	<i>sec</i>	<i>mecA</i>	16s	<i>coa</i>		
ATCC25923, ATCC6538, CMCC26001, 05A046, 05A075, 05A100, 05C209, 05D361, 05F104, 05G021, 05G064, 05G251, 05H150, 05I043, 05L022, 05L216, 05L228, 05N119, 05F318, 05I052, 05L201, 05E162, 05E186, 05E209, 05E024	–	–	–	–	+	+	S	27
ATCC29213, 05B085, 05A188, 05B043, 05C224, 05G226, 05H048, 05I073, 05K037, 05O261, 05O263	+	–	–	–	+	+	S	11
05D004, 05M146	+	–	+	–	+	+	S	2
05L085	+	–	+	+	+	+	S	1
05I056, 05N074, 05L198	–	+	–	–	+	+	S	3
05O114	–	+	+	–	+	+	S	1
SA6	–	–	+	–	+	+	S	1
05C212, 05C215, 05C288, 05C299, 05O073, 05Z073	–	–	–	+	+	+	R	6
05B008, 05B012, 05B033, 05B038, 05B041, 05B059, 05B086, 05B255, 05B289, 05C052, 05C077, 05C078, 05C136, 05C155, 05D131, 05F148, 05F158, 05H058, 05K241, 05P157	+	–	–	+	+	+	R	20
05D154	–	–	+	+	+	+	R	1
Totals	34	4	6	28	71	71	R = 27	

*The tests of methicillin resistance for the bacterial strains were carried out at the Institute of Clinical Pharmacology at Peking University, using the standardized agar dilution method (16, 8, 4, 2 or 1 µg/mL of methicillin, respectively).

S, methicillin sensitive; R, methicillin resistant (a minimal inhibitory concentration greater than 4 µg/mL of methicillin was considered resistant).




FIG. 3. MULTIPLEX POLYMERASE CHAIN REACTION (PCR) OF *STAPHYLOCOCCUS HAEMOLYTICUS* AND *STAPHYLOCOCCUS EPIDERMIDIS* STRAINS

Lane 1, *S. haemolyticus* 05G86; lane 2, *S. epidermidis* 05I078; lane 3, *S. haemolyticus* 05K203; lane 4, *S. haemolyticus* 05G218; lane 5, *Escherichia coli*; lane 6, distilled water (negative control); lane 7, 100 bp DNA marker.

The reliability of the multiplex PCR method was also tested by carrying out the PCR using *S. aureus* DNA in the presence of equal amounts of genomic DNA from other bacteria. There were no unexpected bands when *S. aureus* DNA was mixed with DNA from other bacteria including *Salmonella*, *E. coli*, *L. monocytogenes* and *V. parahaemolyticus* (data not shown). This indicated that the presence of DNA from non-staphylococci strains did not affect the accuracy of the multiplex PCR method.

Sensitivity Test

As shown in Fig. 4, the detection limit of 16S rRNA, *sea* and *mecA* genes was 104.5 pg per reaction (Fig. 4, lane 3), and the detection limit of *coa*, *seb* and *sec* genes was 10.4, 11.1 and 27.6 pg per reaction, respectively (Fig. 4, lanes 3, 11 and 18). Therefore, all the target genes were detectable at 104.5 pg per reaction by the multiplex PCR method.

In the present study, 10^8 cfu of *S. aureus* typically yielded 4.4 µg of DNA (data not shown). This result was similar to that reported by Cremonesi




FIG. 4. SENSITIVITY OF MULTIPLEX POLYMERASE CHAIN REACTION (PCR) ASSAY. Lanes 1–5, quantities of *Staphylococcus aureus* 05B038 DNA were 1.045×10^4 , 1.045×10^3 , 1.045×10^2 , 1.045×10 and 1.045 pg, respectively; lanes 8–12, quantities of *S. aureus* 05L198 DNA were 1.105×10^4 , 1.105×10^3 , 1.105×10^2 , 1.105×10 and 1.105 pg; lanes 15–19, quantities of *S. aureus* SA6 DNA were 2.757×10^4 , 2.757×10^3 , 2.757×10^2 , 2.757×10 and 2.757 pg; lanes 6, 13 and 20, distilled water (negative controls); lanes 7 and 14, 100 bp DNA marker.

et al. (2005), who obtained approximately $10 \mu\text{g}$ of DNA from 10^8 cfu of *S. aureus*. Therefore, the detection limitation can be calculated as 2.4×10^3 cfu. According to the calculation of Yang *et al.* (2002) that 2.78 fg DNA is equivalent to one *S. aureus* genome, the limit of detection of 104.5 pg of the multiplex PCR would be equivalent to 3.64×10^4 genome equivalents. It is well known that *Staphylococcus* spp. grow in clusters of cells, and it has been reported that these clusters of 10–20 cells represent a single cfu (Wilkins *et al.* 1972; Duguid 1989). This means that one cfu of *S. aureus* may actually represent 10–20 genome equivalents. This could explain the apparent 15-fold difference in PCR sensitivity when it is expressed as cfu or genome equivalents.

DISCUSSION

An accurate multiplex PCR method was developed for detecting *S. aureus* and determining the presence of methicillin resistance and enterotoxin genes. This method was proven to be specific and reliable using 71 strains of *S. aureus* and 51 strains of six other bacterial species.

In the multiplex PCR, the detection of *S. aureus* at the genus and species levels was accomplished through the amplification of the genes encoding 16S rRNA and coagulase, respectively. As demonstrated by our multiplex PCR, the primer pair performed in the specific amplification of the 16S rRNA gene from species of *Staphylococcus* and the *coa* primers designed for this study were specific for *S. aureus* strains (Table 3).

It was found that 38.0% (27/71) of the *S. aureus* strains tested were resistant to methicillin. Other studies on clinical *S. aureus* strains detected a similar incidence of MRSA strains (Bach *et al.* 2007); however, there was a lower incidence of methicillin resistance in *S. aureus* isolated from foods (around 25%) (Nitzsche *et al.* 2007). We found a high correlation between methicillin resistance and the presence of the *mecA* gene in the *S. aureus* strains tested. All the MRSA strains yielded *mecA* amplicons, but one *mecA*⁺ strain (*S. aureus* 05L085) was not resistant to methicillin. Furthermore, *mecA* amplicons were identified in some strains of *S. haemolyticus* and *S. epidermidis* in this work. The presence of *mecA* in species of *Staphylococcus* other than *S. aureus* was previously demonstrated (Carneiro *et al.* 2004).

It was found that 77.8% (21/27) of the MRSA strains contained one or more SE gene(s) (*sea*, *seb* or *sec*); however, only 43.2% (19/44) of MSSA strains contained SE gene(s). The reason for the higher incidence of SE genes in MRSA than in MSSA is not thoroughly understood. It has also been reported that the incidence of MRSA that produced SEs was greater than the incidence that did not produce SEs (Moon *et al.* 2007). The presence of virulence genes (including enterotoxin genes) and antibiotic resistance confers a competitive advantage upon *S. aureus* (Hershow *et al.* 1992; Mizobuchi *et al.* 1994), and studies indicate an increase in the isolation of toxin-containing MRSA strains (Takesue *et al.* 1993). Recent studies pointed out that the *mecA* gene is present on the mobile staphylococcal chromosomal cassette *mec* (Katayama *et al.* 2003; Jansen *et al.* 2006), and some of the enterotoxin genes, along with other virulence factors, are part of mobile pathogenicity islands (Novick 2003; Novick and Subedi 2007). Thus, enterotoxin-producing MRSA can develop via the acquisition of *mecA* by toxin-producing MSSA or via the acquisition of toxin genes by MRSA (Layer *et al.* 2006). A recent study indicated that high mobility of the *mecA* gene may be more prevalent than the movement of the enterotoxin genes (Witte *et al.* 2000).

In conclusion, a specific and reliable multiplex PCR method was developed for the detection of 16S rRNA, *coa*, *sea*, *seb*, *sec* and *mecA* genes in *S. aureus* in this study. This is the first PCR method developed for the detection of *S. aureus* that also determines the presence of the genes for methicillin resistance and the foodborne outbreak-associated enterotoxins A, B and C. When coupled with appropriate enrichment and separation procedures, this PCR method will be a useful tool for the detection and identification of *S. aureus* from foods, clinical samples and environmental surveys. In particular, the routine use of this multiplex PCR method for the detection of foodborne *S. aureus* could be used to monitor the presence of enterotoxins and the emergence of methicillin resistance in a population that, to date, has had a relatively low incidence of methicillin resistance.

NOMENCLATURE

MRSA	methicillin-resistant <i>S. aureus</i>
MSSA	methicillin-sensitive <i>S. aureus</i>
SE	staphylococcal enterotoxin

ACKNOWLEDGMENTS

This research was jointly supported by Grant No. 2006BAK02A14 from the Ministry of Science and Technology of China, and Grant Nos. 071422011 and 07dz19508 from the Science and Technology Commission of Shanghai municipality.

REFERENCES

- AKINEDEN, O., ANNEMULLER, C., HASSAN, A.A., LAMMLER, C., WOLTER, W. and ZSCHOCK, M. 2001. Toxin genes and other characteristics of *Staphylococcus aureus* isolates from milk of cows with mastitis. *Clin. Diagn. Lab. Immunol.* 8, 959–964.
- BACH, H.G., STEFFIN, B., CHHADIA, A.M., KOVACHEVICH, R. and GONZALEZ, M.H. 2007. Community-associated methicillin-resistant *Staphylococcus aureus* hand infections in an urban setting. *J. Hand Surg. [Am]* 32, 380–383.
- BALABAN, N. and RASOOLY, A. 2000. Staphylococcal enterotoxins. *Int. J. Food Microbiol.* 61, 1–10.
- BARON, F., COCHET, M.F., PELLERIN, J.L., BEN ZAKOUR, N., LEBON, A., NAVARRO, A., PROUDY, I., LE LOIR, Y. and GAUTIER, M. 2004. Development of a PCR test to differentiate between *Staphylococcus aureus* and *Staphylococcus intermedius*. *J. Food Prot.* 67, 2302–2305.
- CAGATAY, A.A., OZCAN, P.E., GULEC, L., INCE, N., TUGRUL, S., OZSUT, H., CAKAR, N., ESEN, F., ERAKSOY, H. and CALANGU, S. 2007. Risk factors for mortality of nosocomial bacteraemia in intensive care units. *Med. Princ. Pract.* 16, 187–192.
- CARNEIRO, L.A., QUEIROZ, M.L. and MERQUIOR, V.L. 2004. Antimicrobial-resistance and enterotoxin-encoding genes among staphylococci isolated from expressed human breast milk. *J. Med. Microbiol.* 53, 761–768.
- CARROLL, K.C., LEONARD, R.B., NEWCOMB-GAYMAN, P.L. and HILLYARD, D.R. 1996. Rapid detection of the staphylococcal *mecA* gene from BACTEC blood culture bottles by the polymerase chain reaction. *Am. J. Clin. Pathol.* 106, 600–605.

- CHIOU, C.S., WEI, H.L. and YANG, L.C. 2000. Comparison of pulsed-field gel electrophoresis and coagulase gene restriction profile analysis techniques in the molecular typing of *Staphylococcus aureus*. *J. Clin. Microbiol.* **38**, 2186–2190.
- CREMONESI, P., LUZZANA, M., BRASCA, M., MORANDI, S., LODI, R., VIMERCATI, C., AGNELLINI, D., CARAMENTI, G., MORONI, P. and CASTIGLIONI, B. 2005. Development of a multiplex PCR assay for the identification of *Staphylococcus aureus* enterotoxigenic strains isolated from milk and dairy products. *Mol. Cell. Probes* **19**, 299–305.
- DUGUID, J. 1989. *Staphylococcus*: cluster farming gram positive cocci. In *Practical Medical Microbiology*, 13th Ed. (Collee, J.G., Duguid, J.P., Fraser, A.G. and Marmion, B.P., eds.), pp 305–308, Churchill Livingstone, Edinburgh, London, U.K., Melbourne, Australia, New York, NY.
- FOSTER, T.J. 2005. Immune evasion by staphylococci. *Nat. Rev. Microbiol.* **3**, 948–958.
- FRANCOIS, P., PITTET, D., BENTO, M., PEPEY, B., VAUDAUX, P., LEW, D. and SCHRENZEL, J. 2003. Rapid detection of methicillin-resistant *Staphylococcus aureus* directly from sterile or nonsterile clinical samples by a new molecular assay. *J. Clin. Microbiol.* **41**, 254–260.
- HERSHOW, R.C., KHAYR, W.F. and SMITH, N.L. 1992. A comparison of clinical virulence of nosocomially acquired methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* infections in a university hospital. *Infect. Control Hosp. Epidemiol.* **13**, 587–593.
- JANSEN, W.T., BEITSMA, M.M., KOEMAN, C.J., VAN WAMEL, W.J., VERHOEF, J. and FLUIT, A.C. 2006. Novel mobile variants of staphylococcal cassette chromosome mec in *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **50**, 2072–2078.
- JONAS, D., SPECK, M., DASCHNER, F.D. and GRUNDMANN, H. 2002. Rapid PCR-based identification of methicillin-resistant *Staphylococcus aureus* from screening swabs. *J. Clin. Microbiol.* **40**, 1821–1823.
- JORGENSEN, H.J., MORK, T., HOGASEN, H.R. and RORVIK, L.M. 2005. Enterotoxigenic *Staphylococcus aureus* in bulk milk in Norway. *J. Appl. Microbiol.* **99**, 158–166.
- KALIA, A., RATTAN, A. and CHOPRA, P. 1999. A method for extraction of high-quality and high-quantity genomic DNA generally applicable to pathogenic bacteria. *Anal. Biochem.* **275**, 1–5.
- KARAHAN, M. and CETINKAYA, B. 2006. Coagulase gene polymorphisms detected by PCR in *Staphylococcus aureus* isolated from subclinical bovine mastitis in Turkey. *Vet. J.* **174**, 428–431.
- KATAYAMA, Y., ZHANG, H.Z., HONG, D. and CHAMBERS, H.F. 2003. Jumping the barrier to beta-lactam resistance in *Staphylococcus aureus*. *J. Bacteriol.* **185**, 5465–5472.

- LAYER, F., GHEBREMEDHIN, B., KONIG, W. and KONIG, B. 2006. Heterogeneity of methicillin-susceptible *Staphylococcus aureus* strains at a German University Hospital implicates the circulating-strain pool as a potential source of emerging methicillin-resistant *S. aureus* clones. *J. Clin. Microbiol.* **44**, 2179–2185.
- LOWY, F.D. 1998. *Staphylococcus aureus* infections. *N. Engl. J. Med.* **339**, 520–532.
- MASON, W.J., BLEVINS, J.S., BEENKEN, K., WIBOWO, N., OJHA, N. and SMELTZER, M.S. 2001. Multiplex PCR protocol for the diagnosis of staphylococcal infection. *J. Clin. Microbiol.* **39**, 3332–3338.
- MASSEY, R.C., HORSBURGH, M.J., LINA, G., HOOK, M. and RECKER, M. 2006. The evolution and maintenance of virulence in *Staphylococcus aureus*: A role for host-to-host transmission? *Nat. Rev. Microbiol.* **4**, 953–958.
- MCLAUCHLIN, J., NARAYANAN, G.L., MITHANI, V. and O'NEILL, G. 2000. The detection of enterotoxins and toxic shock syndrome toxin genes in *Staphylococcus aureus* by polymerase chain reaction. *J. Food Prot.* **63**, 479–488.
- MEHROTRA, M., WANG, G. and JOHNSON, W.M. 2000. Multiplex PCR for detection of genes for *Staphylococcus aureus* enterotoxins, exfoliative toxins, toxic shock syndrome toxin 1, and methicillin resistance. *J. Clin. Microbiol.* **38**, 1032–1035.
- MIZOBUCHI, S., MINAMI, J., JIN, F., MATSUSHITA, O. and OKABE, A. 1994. Comparison of the virulence of methicillin-resistant and methicillin-sensitive *Staphylococcus aureus*. *Microbiol. Immunol.* **38**, 599–605.
- MOON, J.S., LEE, A.R., KANG, H.M., LEE, E.S., JOO, Y.S., PARK, Y.H., KIM, M.N. and KOO, H.C. 2007. Antibigram and coagulase diversity in staphylococcal enterotoxin-producing *Staphylococcus aureus* from bovine mastitis. *J. Dairy Sci.* **90**, 1716–1724.
- NITZSCHE, S., ZWEIFEL, C. and STEPHAN, R. 2007. Phenotypic and genotypic traits of *Staphylococcus aureus* strains isolated from pig carcasses. *Vet. Microbiol.* **120**, 292–299.
- NOVICK, R.P. 2003. Mobile genetic elements and bacterial toxinoses: The superantigen-encoding pathogenicity islands of *Staphylococcus aureus*. *Plasmid* **49**, 93–105.
- NOVICK, R.P. and SUBEDI, A. 2007. The SaPIs: Mobile pathogenicity islands of *Staphylococcus*. *Chem. Immunol. Allergy* **93**, 42–57.
- OLIVEIRA, D.C., TOMASZ, A. and DE LENCASTRE, H. 2002. Secrets of success of a human pathogen: Molecular evolution of pandemic clones of methicillin-resistant *Staphylococcus aureus*. *Lancet Infect. Dis.* **2**, 180–189.

- OMOE, K., ISHIKAWA, M., SHIMODA, Y., HU, D.L., UEDA, S. and SHINAGAWA, K. 2002. Detection of *seg*, *seh*, and *sei* genes in *Staphylococcus aureus* isolates and determination of the enterotoxin productivities of *S. aureus* isolates Harboring *seg*, *seh*, or *sei* genes. *J. Clin. Microbiol.* **40**, 857–862.
- PADMAPRIYA, B.P., RAMESH, A., CHANDRASHEKAR, A. and VARADARAJ, M.C. 2003. Staphylococcal accessory gene regulator (*sar*) as a signature gene to detect enterotoxigenic staphylococci. *J. Appl. Microbiol.* **95**, 974–981.
- PANLILIO, A.L., CULVER, D.H., GAYNES, R.P., BANERJEE, S., HENDERSON, T.S., TOLSON, J.S. and MARTONE, W.J. 1992. Methicillin-resistant *Staphylococcus aureus* in U.S. hospitals, 1975–1991. *Infect. Control Hosp. Epidemiol.* **13**, 582–586.
- PEREZ-ROTH, E., CLAVERIE-MARTIN, F., VILLAR, J. and MENDEZ-ALVAREZ, S. 2001. Multiplex PCR for simultaneous identification of *Staphylococcus aureus* and detection of methicillin and mupirocin resistance. *J. Clin. Microbiol.* **39**, 4037–4041.
- SABAT, A., MALACHOWA, N., MIEDZOBRODZKI, J. and HRYNIEWICZ, W. 2006. Comparison of PCR-based methods for typing *Staphylococcus aureus* isolates. *J. Clin. Microbiol.* **44**, 3804–3807.
- SHARMA, N.K., REES, C.E. and DODD, C.E. 2000. Development of a single-reaction multiplex PCR toxin typing assay for *Staphylococcus aureus* strains. *Appl. Environ. Microbiol.* **66**, 1347–1353.
- TAKESUE, Y., YOKOYAMA, T., KODAMA, T., SANTOU, T., NAKAMITSU, A., MURAKAMI, Y., IMAMURA, Y., MIYAMOTO, K., OKITA, M., TSUMURA, H., ET AL. 1993. A study on postoperative enteritis caused by methicillin-resistant *Staphylococcus aureus*. *Surg. Today* **23**, 4–8.
- TANG, J.-N., SHI, X.-M., SHI, C.-L. and CHEN, H.-C. 2006. Characterization of a duplex polymerase chain reaction assay for the detection of enterotoxigenic strains of *Staphylococcus aureus*. *J. Rapid Meth. Auto. Microbiol.* **10**, 201–217.
- THOMAS, D.Y., JARRAUD, S., LEMERCIER, B., COZON, G., ECHASSERIEAU, K., ETIENNE, J., GOUGEON, M.L., LINA, G. and VANDEN-ESCH, F. 2006. Staphylococcal enterotoxin-like toxins U2 and V, two new staphylococcal superantigens arising from recombination within the enterotoxin gene cluster. *Infect. Immun.* **74**, 4724–4734.
- UBUKATA, K., NONOGUCHI, R., SONG, M.D., MATSUHASHI, M. and KONNO, M. 1990. Homology of *mecA* gene in methicillin-resistant *Staphylococcus haemolyticus* and *Staphylococcus simulans* to that of *Staphylococcus aureus*. *Antimicrob. Agents Chemother.* **34**, 170–172.

- VIMERCATI, C., CREMONESI, P., CASTIGLIONI, B., PISONI, G., BOET-
TCHER, P.J., STELLA, A., VICENZONI, G. and MORONI, P. 2006.
Molecular typing of *Staphylococcus aureus* isolated from cows, goats and
sheep with intramammary infections on the basis of gene polymorphisms
and toxins genes. J. Vet. Med. B Infect. Dis. Vet. Public Health 53,
423–428.
- WEIGEL, L.M., CLEWELL, D.B., GILL, S.R., CLARK, N.C., MCDOU-
GAL, L.K., FLANNAGAN, S.E., KOLONAY, J.F., SHETTY, J.,
KILLGORE, G.E. and TENOVER, F.C. 2003. Genetic analysis of a
high-level vancomycin-resistant isolate of *Staphylococcus aureus*.
Science 302, 1569–1571.
- WIKLER, M.A., COCKERILL, F.R., CRAIG, W.A., DUDLEY, M.N.,
ELIOPOULOS, G.M., HECHT, D.W., HINDLER, J.F., LOW, D.E.,
SHEEHAN, D.J., TENOVER, F.C., ET AL. 2006. *Methods for Dilution
Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically;
Approved Standard*. Clinical Laboratory Standards Institute, Wayne, PA.
- WILKINS, J.R., DARNELL, W.L. and BOYKIN, E.H. 1972. Cinemicro-
graphic study of the development of subsurface colonies of *Staphylococ-
cus aureus* in soft agar. Appl. Microbiol. 24, 786–797.
- WITTE, W., BRAULKE, C., HEUCK, D. and CUNY, C. 2000. Methicillin
resistant *Staphylococcus aureus* in German hospitals develop narrower
patterns of antimicrobial resistance. Euro Surveill. 5, 31–34.
- YANG, S., LIN, S., KELEN, G.D., QUINN, T.C., DICK, J.D., GAYDOS,
C.A. and ROTHMAN, R.E. 2002. Quantitative multiprobe PCR assay for
simultaneous detection and identification to species level of bacterial
pathogens. J. Clin. Microbiol. 40, 3449–3454.
- ZHANG, K., SPARLING, J., CHOW, B.L., ELSAYED, S., HUSSAIN, Z.,
CHURCH, D.L., GREGSON, D.B., LOUIE, T. and CONLY, J.M. 2004.
New quadriplex PCR assay for detection of methicillin and mupirocin
resistance and simultaneous discrimination of *Staphylococcus aureus*
from coagulase-negative staphylococci. J. Clin. Microbiol. 42, 4947–
4955.